




Amendments to the Claims

1-43. (Cancelled)

44. (Currently amended) A metallocene-based ligand having a formula selected from the group consisting of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V), Formula (VI), Formula (VII), Formula (VIII), and Formula (IX):



wherein

W is phosphorus or arsenic;

M is a metal;


R^1 and R^2 are different from each other and are independently selected from the group consisting of unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, unsubstituted heteroarylamino, substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, and substituted heteroarylamino;

R^3 and R^4 are independently selected from the group consisting of substituted branched-chain alkyl, substituted straight-chain alkyl, substituted cycloalkyl, substituted carbocyclic aryl, substituted heteroaryl, unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted cycloalkyl, unsubstituted carbocyclic aryl, and unsubstituted heteroaryl;

n is an integer from 0 to 3;

m is an integer from 0 to 5;


Q is the group



wherein

R^8 is selected from the group consisting of substituted straight-chain alkyl, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl; R^9 and $R^{10''}$ are independently selected from the group consisting of hydrogen, substituted straight-chain alkyl, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl; or

Q is selected from the group consisting of



wherein


R^6 and R^7 are independently selected from the group consisting of substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, substituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, substituted heteroarylamino, unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, and unsubstituted heteroarylamino;

R^8 , R^9 , R^{10} and $R^{10''}$ are independently selected from the group consisting of hydrogen, substituted straight-chain alkyl, unsubstituted straight-chain alkyl, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl; R^{11} is selected from the group consisting of OR^{13} , SR^{13} , NHR^{13} , and $NR^{13}R^{14}$, wherein

R^{13} and R^{14} are independently selected from the group consisting substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and

unsubstituted heteroaryl; R^{12} is selected from the group consisting of hydrogen, halogen, OR^{13} , SR^{13} , $NR^{13}R^{14}$, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl, and n' is 0 to 4;


R^5 is selected from:



wherein R^{15} , R^{16} and R^{17} are independently selected from the group consisting of hydrogen, ~~halogen~~ halogen, OR^{13} , SR^{13} , $NR^{13}R^{14}$, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl substituted carbocyclic aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl; and wherein the two geminal substituents R^{18} together are a doubly bonded oxygen atom, or each substituent R^{18} is individually hydrogen; and G is selected from the group consisting of $-C(=O)NH-R^*$, $-NHCO-$, $-C(=O)-OR^*O-C(=O)-$, $-C(=O)-R^*C(=O)-$, $-CH=N-R^*-N=CH-$, $-CH_2NH-R^*-NHCH_2-$, $-CH_2NHC(=O)-R^*-C(=O)NHCH_2-$, $-CH(R^8)NH-R^*-NH(CH(R^8)-$, $-CH(R^8)NHC(=O)-R^*-C(=O)NHCH(R^8)-$, $-C(=O)NH-R-NHC(=O)-$, $-C(=O)-ORO-C(=O)-$, $-C(=O)-RC(=O)-$, $-CH=N-R-N=CH-$, $-CH_2NH-R-NHCH_2-$, $-CH_2NHC(=O)-R-C(=O)NHCH_2-$, $-CH(R^8)NH-R-NH(CH(R^8)-$, $-CH(R^8)NHC(=O)-R-C(=O)NHCH(R^8)-$;

wherein R^8 is, independently, as previously defined;

$-R^*$ - and $-R-$ are selected from the group consisting of:




wherein R^{12} is as previously defined;

wherein the two substituents R^{19} together are $-(CH_2)_m-$ or each substituent R^{18} is independently selected from the group consisting of hydrogen, substituted branched-chain alkyl, unsubstituted branched-chain alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted carbocyclic

aryl, unsubstituted carbocyclic aryl, substituted heteroaryl, and unsubstituted heteroaryl; wherein the or each heteroatom is independently selected from sulphur, nitrogen, n' is an integer of from 0 to 4; and m' is an integer of from 1 to ~~8~~; 8

~~with the proviso that the following compounds are excluded:~~




45. (Previously presented) The metallocene-based ligand of Claim 44, which is a diastereomer having Formula (IV), Formula (V), or Formula (VI).

46. (Previously presented) The metallocene-based ligand of Claim 44, which is an enantiomer having Formula (VII), Formula (VIII), or Formula (IX).

47. (Previously presented) The metallocene-based ligand of Claim 44, wherein the metallocene-based ligand is a phosphine or arsine having chirality at W, and wherein the metallocene-based ligand has at least one additional element of chirality selected from the group consisting of chirality at carbon, and axial chirality.

48. (Previously presented) The metallocene-based ligand of Claim 44, wherein the metallocene-based ligand is a diphosphine or diarsine having chirality at W, and wherein the metallocene-based ligand has two additional elements of chirality comprising chirality at carbon, and axial chirality.

49. (Previously presented) The metallocene-based ligand of Claim 44, wherein the metallocene is ferrocene.
50. (Previously presented) The metallocene-based ligand of Claim 44, wherein W is phosphorus.
51. (Previously presented) A catalyst or catalyst precursor in an asymmetric transformation reaction to generate a high enantiomeric excess of a formed compound, the catalyst or catalyst precursor comprising the metallocene-based ligand of Claim 44.
52. (Previously presented) A transition metal complex containing a transition metal coordinated to a ligand according to the metallocene-based ligand of Claim 44.
53. (Previously presented) A transition metal complex according to claim 52, wherein the transition metal is a Group VIb or a Group VIII metal.
54. (Previously presented) A method for preparing the metallocene-based ligand of Claim 44, comprising:
providing a metallocene-based substrate having a chiral directing substituent on one or both rings;
ortho-lithiating the metallocene-based substrate; and
converting the ortho-lithiated metallocene-based substrate to obtain the metallocene-based ligand.
55. (Previously presented) The method according to Claim 54, wherein the metallocene-based ligand has Formula (I) or Formula (III), wherein the metallocene-based substrate has Formula (X'):




Formula (X')

wherein R^3 and R^4 are independently selected from the group consisting of substituted branched-chain alkyl, substituted straight-chain alkyl, substituted cycloalkyl, substituted carbocyclic aryl, substituted heteroaryl, unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted cycloalkyl, unsubstituted carbocyclic aryl, and unsubstituted heteroaryl;

n is an integer from 0 to 3;

and wherein X^* is a chiral directing group, wherein the step of converting the ortho-lithiated metallocene-based substrate comprises reacting the ortho-lithiated substrate with an R^1 substituted phosphine or arsine, and with an R^2 -bearing Grignard reagent or an R^2 -organolithium compound, then converting X^* to Q or G.

56. (Previously presented) A method according to Claim 55, wherein X^* is selected from the group consisting of:



wherein

R^a and R^b are independently selected from the group consisting of substituted branched-chain alkyl, substituted straight-chain alkyl, substituted cycloalkyl, substituted carbocyclic aryl, substituted heteroaryl, unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted cycloalkyl, unsubstituted carbocyclic aryl, and unsubstituted heteroaryl.

57. (Previously presented) The method according to claim 55, wherein the ortho-lithiation step is conducted using at least one lithiating agent selected from the group consisting of n-butyllithium, sec-butyllithium, and tert-butyllithium.


58. (Previously presented) The method according to claim 57, wherein the step of converting the ortho-lithiated metallocene-based substrate comprises reacting the ortho-lithiated metallocene-based substrate *in situ* with a dichlorophosphine of the formula R^1PCl_2 wherein R^1 is selected from the group consisting of unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, unsubstituted heteroarylamino, substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, substituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, and substituted heteroarylamino;

to yield an intermediate product, wherein the intermediate product is converted to obtain the metallocene-based ligand.

59. (Previously presented) The method according to Claim 58, further comprising reacting the intermediate product with an organometallic reagent of formula R^2Z , wherein R^2 is selected from the group consisting of unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted

carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, unsubstituted heteroarylamino, substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, substituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, and substituted heteroarylamino;

wherein Z is Li or MgY, and wherein Y is a halide, to obtain a phosphorus chiral compound having formula (XI'):




Formula (XI')

wherein the phosphorous chiral compound is converted to obtain the metallocene-based ligand.

60. (Previously presented) The method of Claim 59, wherein the metallocene-based ligand has Formula (I) or Formula (III).

61. (Previously presented) A method for preparing a metallocene-based ligand of Claim 44, comprising:

providing a compound of Formula (XXXVII):




wherein X is an achiral directing group;

subjecting the compound of Formula (XXXVII) to enantioselective mono-ortho-lithiation using


at least one lithiating agent selected from the group consisting of n-butyllithium, sec-butyllithium, and tert-butyllithium, wherein the mono-ortho-lithiation is conducted in the presence of a homochiral tertiary amine, whereby a chiral monolithium compound is obtained; reacting the chiral monolithium compound *in situ* with a dichlorophosphine of the formula R^1PCl_2 followed by reacting with an organometallic reagent of the formula R^2Z , wherein R^1 and R^2 are different from each other and are independently selected from the group consisting of unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, unsubstituted heteroarylamino, substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, substituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, and substituted heteroarylamino;

wherein Z is Li or MgY, and wherein Y is a halide, to obtain a phosphorus chiral compound having Formula (XXXVIII):



and converting the phosphorus chiral compound having Formula (XXXVIII) to the metallocene-based ligand, wherein the metallocene-based ligand has Formula (I) or Formula (III).


62. (Previously presented) The method according to Claim 61, wherein X is selected from the group consisting of:



wherein R^a and R^b are independently selected from the group consisting of substituted branched-chain alkyl, substituted straight-chain alkyl, substituted cycloalkyl, substituted carbocyclic aryl, substituted heteroaryl, unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted cycloalkyl, unsubstituted carbocyclic aryl, and unsubstituted heteroaryl.

63. (Previously presented) A method for preparing a metallocene-based ligand of Claim 44, comprising:


providing a compound of the Formula (XXXIX):



wherein X^* is a chiral directing group;

subjecting the compound of Formula (XXXIX) to bis-ortho-lithiation using at least one lithiating agent selected from the group consisting of n-butyllithium, sec-butyllithium, and tert-butyllithium, whereby a bislithium compound *in situ* with a dichlorophosphine of the formula R^1PCl_2 followed by reacting with an organometallic reagent of the formula R^2Z wherein R^1 and R^2 are different from each other and are independently selected from the group consisting of unsubstituted branched-chain alkyl, unsubstituted straight-chain alkyl, unsubstituted alkoxy, unsubstituted alkylamino, unsubstituted cycloalkyl, unsubstituted cycloalkoxy, unsubstituted cycloalkylamino, unsubstituted carbocyclic aryl, unsubstituted carbocyclic aryloxy, unsubstituted heteroaryl, unsubstituted heteroaryloxy, unsubstituted carbocyclic arylamino, unsubstituted heteroarylamino, substituted branched-chain alkyl, substituted straight-chain alkyl, substituted alkoxy, substituted alkylamino, substituted cycloalkyl, substituted cycloalkoxy, substituted cycloalkylamino, substituted carbocyclic aryl, substituted carbocyclic aryloxy, substituted heteroaryl, substituted heteroaryloxy, substituted carbocyclic arylamino, and substituted heteroarylamino;

wherein Z is Li or MgY , and wherein Y is a halide, to obtain a phosphorus chiral compound having Formula (XXXX):



and converting the phosphorous chiral compound having Formula (XXXX) to the metallocene-based ligand, wherein the metallocene-based ligand has Formula (II).